

Placement of dental implants usually does not create a great deal of pain. General post-operative instructions apply, with the exception of rinsing. Remember, **DO NOT** rinse with the warm salt water. The doctor has given you a prescription rinse containing chlorhexidine, an antibacterial medicine, to assist in the healing and keep the site healthy. Make sure that you get rid of any toothpaste residue prior to rinsing. **DO NOT** eat or drink for 30 minutes after rinsing. This will maximize the potency of the medication.

Generally speaking, following implant surgery, the Gore-Tex sutures need to be removed. They are white strings which may see in your mouth. If they become loose and annoying, call the office to have them removed at your earliest convenience.

1. Some degree of **DISCOMFORT AND PAIN** arises as numbness subsides. At the first sign of pain or discomfort, take the prescribed medications. As an alternative, you may take 2-4 Advil tablets. If you cannot take aspirin products, then take two Tylenol tablets. If the pain you are having is not controlled by the Advil, or "Advil-like" prescription, you may take the prescribed severe pain medicine (typically Vicodin ES), but only after eating some food. Any pain medications can cause nausea and vomiting. It is very important that you have some food in your stomach before you take them.

2. **DO NOT DISTURB THE AREA OF SURGERY.** Allowing the tissues to rest undisturbed assists the healing process. Avoid vigorous chewing, excessive spitting, or aggressive rinsing. If you routinely use a Water-Pik, avoid doing so during the healing phase. Initial healing may be delayed, active bleeding restarted, or infection introduced.

3. **EXPECT MINOR BLEEDING OR OOZING.** If bleeding persists, continue pressure on a fresh sponge for an additional 30 minutes to an hour. Biting on a moist teabag wrapped in gauze may help control persistent oozing from the surgical site. Tea contains tannic acid, which helps promote blood clotting.

4. **FIRM PRESSURE FOR 15-30 MINUTES** usually controls the problem. If you check too frequently (every few minutes), the blood clot will not form properly and bleeding will continue. Should active bleeding persist, please call the office.

5. **LIMIT PHYSICAL ACTIVITY** during the first 24-48 hours after surgery. Over exertion may lead to post-operative bleeding and discomfort. When you lie down, keep your head elevated on a pillow. You may wish to place a towel on your pillowcase to avoid staining from any blood-tainted saliva.

6. **PAIN FOLLOWING ORAL SURGERY** will be most severe within the first 6-8 hours after the operation. Please do not drink alcoholic beverages while taking prescription medication. Do not wait for the pain to become unbearable before using some form of pain medication, as then it will be more difficult to control. Moderate to severe pain usually does not last longer than 48-hours. Mild discomfort usually diminishes after the third day. Please contact us if you experience prolonged severe pain so that we can help make you more comfortable.

7. **SWELLING RELATED TO THE SURGICAL PROCEDURE** may occur following certain types of procedures. An ice pack should be placed on the side of your face for 20 minutes then taken off for 10 minutes and repeated. Anti-inflammatory medications, such as Advil also help decrease swelling.

8. **FLUID INTAKE IS IMPORTANT.** Clean beverages, ginger ale, 7-up, or Sprite, water, teas, soda, broth, soups, or juices are all suitable. Avoid hot liquids until the numbness has worn off, and the bleeding has stopped. Drink plenty of fluids.

9. **AVOID USING A STRAW FOR SEVERAL DAYS.** It may dislodge the blood clot and delay healing.

10. **FOOD SELECTION.** Soft, cool, foods are most easily tolerated. A nutritious diet is most important to your comfort and temperament. Remember that eating can prevent nausea sometimes associated with certain medications. Supplements such as Ensure, Carnation Instant Breakfast, and/or yogurt supply excellent added nutrition.

***DO NOT EAT ANY POPCORN, POPPY SEEDS, SESAME SEEDS, OR SEEDED FRUITS. THEY CAN GET STUCK AND CAUSE INFECTION!**

11. Take any **SPECIAL MEDICATION** we have prescribed on the special dosing schedule. yogurt with active cultures or acidophilus should be taken while on antibiotics to prevent diarrhea. It is important to take the antibiotics to completion. If you are given antibiotics and take birth control pills, you should be aware that the birth control may become ineffective, therefore take appropriate precautions.

12. Take any **REGULARLY SCHEDULED MEDICATION** (for diabetes, high blood pressure, etc.) on your regular schedule unless advised to do otherwise.

13. **TRY TO AVOID SMOKING COMPLETELY**, as it tends to delay healing and interfere with regeneration.

14. **THE DAY FOLLOWING SURGERY AND THEREAFTER.** On the day following the surgery, rinse your mouth carefully with 1/2 teaspoon of salt to a large glass of warm water. Repeat three times a day until remaining soreness subsides. Please do not use a syringe or Water-Pik during the healing phase.

15. **DO NOT WORRY ABOUT STICHES.** They are usually dissolvable.

16. **ANY SWELLING, SORENESS, OR STIFFNESS IN THE JAW MUSCLES** can be relieved by applying a warm moist towel to the affected side of the face.

Sometimes a soft diet may be necessary for the first few days following surgery. Most patients are able to resume regular food intake within a short time.

Bruising marks may appear on the skin of the face during the first few days after surgery. Should any undue reaction or complications arise, notify the office immediately.

If you have any problems or questions please do not hesitate to call our office at (215) 223-5555.

We will make every effort to contact you ASAP

Thank you!